MEN'S HEALTH ASSESSMENT QUESTIONNAIRE

Λ	110	B		е тм
H	VO		WI	100
46	1		W 4 W	-
	Mill	aster	idel	

Name		Date					
American Urological Association Symptom Index (AUA-		the answe	r that best	describes	s your sym	ptoms.	
Urinary symptoms during the past month	Not at all	Less than 1 time in 5	Less than half the time	About half the time	More than half the time	Almost always	
How often have you had a sensation of not emptying your bladder completely?	0	1	2	3	4	5	
2. How often did you urinate more than once within a 2-hour period?	0	1	2	3	4	5	
3. How often have you stopped and started several times while urinating?	0	1	2	3	4	5 /	
4. How often have you had difficulty postponing urination?	0	1	2	3	4	5	
5. How often have you had a weak urinary stream?	0	1	2	3	4	5	
6. How often did you strain to begin to urinate?	0	1	2	3	4	5	
7. How many times did you get up during the night to urinate?	0 TIMES	1 TIME	2 TIMES	3 TIMES	4 TIMES	5 TIMES	
Adapted from the AUA-SI BOTHER ASSESSMENT QUESTION	Total Score:						
Overall, how bothersome has any trouble with urination be \(\sim\) Not at all bothersome \(\sim\) Bothers me a little \(\sim\) E				s me a lot	L		
Sexual Function Question							
Nould you like to talk to the doctor about sexual function?	□ yes □ no						

AVODART is indicated for the treatment of symptomatic benign prostatic hyperplasia (BPH) in men with an enlarged prostate to improve symptoms, reduce the risk of acute urinary retention, and reduce the risk of the need for BPH-related surgery.

Only your doctor can tell if your symptoms are from an enlarged prostate and not a more serious condition such as prostate cancer. See your doctor for regular exams.

Women and children should not take AVODART 0.5mg Soft Gelatin Capsules. Women who are, or could become pregnant, should not handle AVODART due to the potential risk to a developing male fetus.

Men treated with AVODART should not donate blood until at least 6 months after their final dose. Tell your doctor if you have liver disease. AVODART may not be right for you.

Possible side effects include sexual side effects, breast tenderness and/or swelling. These side effects occur infrequently.

